Index to Volume 83 of the Bryologist[1,2]

Acarospora fuscata, 330; lorentzii, 219
Acaulon muticum, 202
achlorophyllous spores, in mosses, 554
acid-base color reactions, in mosses, 228+; the status of Triquetrella ferruginea, Barbula inaequalifolia and B. calcarea, 228
acrocarpous mosses, effects of trampling, 311; rhizoids, 481
Acrocladium auriculatum, 443; chlamydophyllum, 443
Acroporium pungens, 556
adaptive advantage, asexual reproduction, 344
Agyrophora lyngei, 91
air pollution, effects on lichen communities, 281, 296
Alectoria loxensis, 530; nigricans, 90+; ochroleuca, 88+
A. LeRoy Andrews Bryological Foray of 1979, 130; of 1980, 129
algal-fungal relation, in appressorum-type, 177; in Chiodecton sanguineum, 176+
ALLEN, B. H. Mosses new or noteworthy from Colorado, 257
ALLEN, B. H. Peristome variation in the genus Fissidens: an SEM study, 314
Aloinia aloides var. ambigua, capsule as a food source for ants, 524+, 525*
Amazon Basin, western, 526
Amblystegiaceae, 512, 523
Amblystegium juratzkanum, 202, 513; riparium, carotenoids of, 14+; varium, 202, var. alaskanum, 513, var. lancifolium, 513, var. ovatum, 513, var. parvulum, 513, var. varium, 513
amoeboid plastids, 147+; in Anthoceros punctatus, 157; in Isoetes, 147
Amphidium lapponicum, 365
Anacolia menziesii, in Colorado, 253
Anaptychia ulotrichoides, 251+, 252*
Anastrophyllum hellerianum, 43+, 95; minutum, 238, var. minutulum, 95
ANDERSON, L. E. (review), A. J. E. SMITH, The moss flora of Britain and Ireland, 394
Andreaea acutifolia, 212+, 443; in Hawaii, 212; mutabilis, 212, 443; perpapillosa, 212; rupestris, 212+, 238, 443
ANDRUS, R. E. A large range extension for Sphagnum henryense, 239
ANDRUS, R. E. Sphagnum **andersonianum**, a new species endemic to northeastern North America, 60
Aneura pinguis, 95, 352
Anictangium repens, 475
Anisomeridium biforme, 4; **feeanum**, 4; **subprostans**, 4; **tamarindi**, 4; **tuckeri**, 4, 5*
Anisothecium pallidisetum, 560+
Annual meeting, American Bryological and Lichenological Society, 403
Anoectangium aestivum, 229, 444, 556
Anomobryum filiforme, 556
Anomodon attenuatus, 161+, 202; californicus, 230, 363; minor, 164+, 202; rostratus, 164, 202; scaberrimus, 419; viticulosus, 164+
Anthelia julacea, 95; juratzkana, 351+
antheraxanthin, 22+
antheridial development, in Jubula, 194
Anthoceros punctatus, amoeboid plastids in, 157
Anthocyanins, in Pottiaceae, 228
Anthracothecium cf. pyrenuloides, 6; desquamans, 368; hians, 368; leucostomum, 6; oligosporum, 368; thelomorphum, 6
Antitrichia, 455; curtipendula, 358
Aplodon wormskioldii, 238
Apometzgeria pubescens, 95
apothecia, imperforate, 345; in Parmotrema, 345; perforate, 345
appressorum-type, algal-fungal relationship in, 177
archegonial development, in Jubula, 195
archesporial cell, in Ditrichum pallidum, 141
archesporial wall, 138+
Archidium, 536; donnellii, 202; ohioense, 202
Archifissidentaceae, 559
Arctoa fulvella, 361
Arcto-Tertiary Geoflora, 441, 519
Argentina, lichens in, 219
Arthonia caesia, 288+; macounii, 71
Arthopyrenia, 1; sect. Polymeridium, 12; affinis, 18; bifera, 6; cinchonae, 6; contendens, 12; feeana, 4; lyrata, 6; **lyrata**, 8*; **minor**, 5*, 6+; padi, 6+; prospersella, 13; salicis, 6+; subcinerea, 12; subprostrans, 6; **tenuis**, 18; tichothecioides, 13
ascospore, size in relation to longevity, 348
asexual reproduction, energy investment in, 344; genetic costs in, 348
asexual reproductive patterns, in Parmotrema, 344; strategies, adaptive advantage in, 344
Aspicilia, 331; cinera, 330+; gibbosa, 328+
a-taxanthin, 22+
Asterella gracilis, 95; saccata, 95
Astomiopsis, 475; radiculosa, 255; saint-pierrei, 254+; sinensis, 255
Astomum muhlenbergianum, 202; phascoides, 203
Astrothelium pyrenastraeum, 13

[1] Compiled by Paul L. Redfearn, Jr., Department of Life Sciences, Southwest Missouri State University, Springfield, MO 65802.

[2] New scientific names are printed in boldface type; illustrations are indicated by an asterisk; taxa discussed or mentioned several times in a paper are indicated by a plus sign.

atranorin, 331, 341
Atrichum altecristatum, 238; angustatum, 73, 203, 372, chlorophyll a/b ratio, 85, sensitivity to trampling, 301, var. rhystophyllum, 556; undulatum, 203, 542
Aulacomniaceae, 475
Aulacomnium, 90; acuminatum, subfossils, 487; heterostichum, 203; palustre, 203, 492; turgidum, 238, 492
aurochrome, 22+
auroxanthin, 22+
auxin, exogenous effect on Riccia, 67

Bacidia chlorantha, 288+; chlorococca, 288+
Barbella pendula, 556
Barbilophozia barbata, 95; floerkei var. floerkei, 95; hatcheri, 95, 352; kunzeana, 95; lycopodioides, 95, 352; quadriloba, 95
Barbula, 230+; brachyphylla, 356; calcareum, 231+; cancellata, 203; convoluta, 203, 489; ehrenbergii, 203; fallax, 363; inaequalifolia, 231+; indica, 349; orizabensis, 511; platyneura, 489; pringlei, 556; revoluta, 489; subulifolia, 556; torquata, reduced branches in, 483; unguiculata, 203, 238, 536
Bartramia, 280; ithyphylla, 238, 365; pomiformis, 203, 238; stricta, 253
Bauer, C. R. *See* Crosby, M. R. & C. R. Bauer, 603
Bazzania ambigua, 95; denudata, 95, 238; tricrenata, 95; trilobata, 238, 307
Belland, R. J. & G. R. Brassard. A major range extension for Encalypta longicolla, 563
Belland, R. J. & G. R. Brassard. New or additional moss records from Newfoundland VI, 360
Belonia americana, 17
Bermudez R., I. R. *See* Reese, W. D. & I. R. Bermudez R., 250
Bestia, 455
Biatora columbiana, 71
Big Thickett, Sphagnum in, 247; Preserve, bryophytes in, 369
Bilewsky, F. Obituary, 279
bioindicators, 281, 296
biomass, lichens, 32
birds, dispersal of mosses, 493
Bizot, M., notice of death, 129
Blepharostoma arachnoideum, 95; trichophyllum subsp. brevirete, 95; trichophyllum subsp. trichophyllum, 56, 95, 353
Blinda acuta, 238
"bouquet" stage, 143+, 144*
Brachelyma subulatum, 203
Brachymenium bulbiferum, 556
Brachytheciaceae, 512
Brachythecium acuminatum, 164, 203; asperimum, 253; calcareum, 203; curtum, 203, 306; oxycladon, 164+, 203; plumosum, 203; rivulare, 164+; rutubalum, 164+, 203; salebrosum, 203; speciosum, 556; stereopoma, 556; velutinum, chlorophyll a/b ratio, 85
Bragg, T. B. (review), N. G. Slack, Species diversity and community structure in bryophytes, 400
Brassard, G. R. Bryophyta Exsiccata Terrae-Novae et Labradoricae: A new exsiccata from Newfoundland and Labrador, Canada, 237
Brassard, G. R. *See* Belland, R. J. & G. R. Brassard, 360, 563
Brassard, G. R. *See* Fife, A. J. & G. R. Brassard, 358
Bray-Curtis polar ordination, 284+
Breutelia arcuata, 73; jamaicensis, 556; tomentosa, 556
Britton, E. B., 335
Brodo, I. M. (review), R. B. Filson and R. W. Rogers, Lichens of South Australia, 614
Brotherella recurvans, 306+; tenuirostris, 203
Brown, R. C. & B. E. Lemmon. Ultrastructural aspects of chloroplast development in spores of the moss Leptodictyum riparium (Hedw.) Warnst., 545
Brown, R. C. & B. E. Lemmon. Ultrastructure of sporogenesis in a moss, Ditrichum pallidum I. Meiotic prophase, 137
Brown, R. C. & B. E. Lemmon. Ultrastructure of sporogenesis in a moss, Ditrichum pallidum II. Metaphase I through the tetrad, 153
Bruchia bolanderi, 356; drummondii, 545; texana, 203
Bryhnia graminicolor, 163+
Bryoandersonia illecebra, 164+, 203
Bryoerythrophyllum, 230+; binnsii, 367; **calcareum**, 232; **inaequalifolium**, 231; jamesonii, 367, 492; recurvirostrum, 238, 489
bryoides-type peristome, 316
Bryological and Lichenological Section, Botanical Society of America, 129; meeting, 403
Bryological Times, 130
Bryology at the XIIIth International Botanical Congress, 130
Bryomanginia radiculosa, 255; saint-pierrei, 255
bryophyte communities along trails, 302
bryophyte exsiccata from Labrador, 237
bryophytes, calciphilous, 161+; frequency in sinks, 162+; saxicolous, 165; sensitivity to trampling, 301; stress adaptation, 349
bryophytes in sinks, cover, 162+; importance values, 163+; relative cover, 162+; relative frequency, 162+
Bryopteris 181+, 182*
Bryoria chalybeiformis, 91
Bryoxiphium norvegicum, new to Iowa, 535
Bryson, C. T. Diphyscium foliosum new to Mississippi, 372
Bryum, 280, 307, 536; argenteum, 203, 311; bicolor, 203, 356, capsules as food source for ants, 524, 525*; caespiticium, 203; capillare,

164, 203; creberrimum, 203; gemmiparum, 203; limbatum, 556; muehlenbeckii, 203; pseudocapillare, 556; pseudotriquetrum, 492, var. bimum, 203; weigelii, 238
BUCK, W. Animadversions on Pterigynandrum with special commentary on Forsstroemia and Leptopterigynandrum, 451
BUCK, W. R. (review), P. TIXIER, Bryogeographie du Mont Bokor (Cambodge), 277
BUCK, W. R. & R. H. ZANDER. On Astomiopsis, Bryomanginia and Melophyllum (Ditrichaceae), 254
Buellia disciformis, 91, 330, var. muscorum, 89+; punctata, 288; schaereri, 288
Buxbaumia, 280; aphylla, 238

calcareous substrates, 337
calcifuges, 161
calciphiles, 161
calciphilous bryophytes, 161+
Calicium subcurtum, 219
Callicaldium haldanianum, sensitivity to trampling, 301
Calliergon aftonianum, 492+
Caloplaca cinnabarina, 219; constipans, 64; coralloides, 66; discolor, 288+; erythrantha, 219; stillicidorum, 89+; sublobulata, 219; tiroliensis, 89+
Calypogeja fissa, 307+; muelleriana, 96; sphagnicola, 96; suecica, 96; trichomanis, 96
Calyptopogon, 474
Campylium chrysophyllum, 164+, 203; halleri, 254; hispidulum, 492, var. sommerfeltii, 203; radicale, 492
Campylopus, distribution in North America, 584*, 585*, in North America, 570, key to North American taxa, 571; adustus, 572+; angustiretis, 570+, 582; arctocarpus, 570+; arctocarpus, distribution in North America, 584*; arenicola, 582; atrovirens, 238, 570+; atrovirens, distribution in North America, 584*, var. **cucullatifolius**, 571+, 573*, var. cucullatifolius, distribution in North America, 584*; aureus, 571+, 575*, 576, distribution in North America, 585*; bermudianus, 570+, 588; carolinae, 570+, 573*, distribution in North America, 584*; chionophilus, 586; chrysodictyon, 586; cuspidatus, 574; cygneus, 556; delicatulus, 577, var. carolinae, 577; dicnemioides, 574; donnellii, 570+, 582; exasperatus, 574; flexuosus, 570+, 577, distribution in North America, 584*; fragilis, 556, 570+, 578, distribution in North America, 585*; gracilicaulis, 570+, 577, 581+, var. brevicaulis, 581, var. donnellii, 570+; introflexus, 443, 570+, 576, distribution in North America, 584*; japonicus, 574; lamprodictyon, 586; mildei, 579; oerstedianus, 571+, 573*, 579, distribution in North America, 584*; paradoxus, 587; paradoxus, 587; pauper, 586; pilifer, 443, 571+, 579, distribution in North America, 584*; polytrichoides, 578+; porphyreodictyon, 556; richardii, 556; rosulatus, 586; schimperi, 570+, 580, distribution in North America, 585*; schwarzii, 570+, 580, distribution in North America, 585*; subg. Pseudocampylopus, 581; subleucogaster, 570+, 581+; subulatus, 359, 571+, 581, distribution in North America, 585*, var. schimperi, 570+, 580; surinamensis, 570+, 573*, 577, 581+, distribution in North America, 585*, var. **angustiretis**, 571+, 582, var. surinamensis, 571+; tallulensis, 570+, 577, distribution in North America, 585*, var. subleucogaster, 570+; trachyblepharon, 571+, 588, distribution in North America, 585*; trollii, 586
Campylothelium, 7; **amylosporum**, 7, 8*, 9; nitidum, 7; zahlbruckneri, 7
Candelaria concolor, 288, var. effusa, 288; fibrosa, 219
Candelariella athallina, 91; aurella, 91; kuusamoensis, 91; vitellina, 90+; xanthostigma, 288
caperatic acid, 331+
carbon dioxide, compensation levels in lichens, 505; exchange, in lichens, 505
α-carotene, 22+, β-, 22+, γ-, 22+; β-carotene diepoxide, 22+, epoxide, 22+
carotenoids, Bryophyta, 21
Catillaria **columbiana**, 71+; hypochlorella, 72; laureri, 72
Catoscopium nigritum, 492
cell disruption bomb, 342
Cephalozia bicuspidata, 353, subsp. bicuspidata, 95, subsp. lammersiana, 96; connivens var. connivens, 96; leucantha, 96; lunulifolia, 96, 307; pleniceps, 353, var. pleniceps, 96; divaricata, 353, var. divaricata, 96, var. scabra, 96; hyalania, 307; phyllacantha, 225+, ecology, 226; rubella var. rubella, 96
Ceratodon heterophyllus, 234+; purpureus, 203, 234+, 238, 311, carotenoids, 24+, var. **rotundifolius**, 234+, 235*, distribution of, 236*
Cetaria cucullata, 90+; delisei, 89+; islandica, 91; nigricans, 90+; nivalis, 91; oakesiana, 330
Chaenomitrium, 405+
Chamaebryum, 467+; character states in, 472+; pottioides, 475
Chandonanthus setiformis, 238
checklist of mosses of Oklahoma, 202
Chelan Nat'l. Rec. Area, hepaticae of, 95
Chiloscyphus pallescens var. pallescens, 96; polyanthos, 164, 353, var. polyanthos, 96, var. rivularis, 96, 353
Chiodecton sanguineum, 170+; algal-fungal relationship in, 176
chlorophyll a/b ratios, 84
chlorophyllous spores, in mosses, 554
chloroplast development, in Leptodictyum riparium, 545+, 546*–552*; maturation, in Leptodictyum riparium, 545+, 546*–552*; ultra-

structure in Leptodictyum riparium, 545+, 546*–552*
CHRISTY, J. A. Additions to the moss flora of Oregon, 355
CHRISTY, J. A. Rediscovery of Sciaromium tricostatum (Sull.) Mitt. (=Limbella tricosta) in Oregon, 521
Cirriphyllum piliferum, 164
Cladonia amaurocraea, 90+; chlorophaea, 91; coccifera, 91; coniocraea, 288; cristatella, 176; ecmocyma, 90; furcata, 330; gracilis, 89+; mitis, 90+; pleurota, 92; pocillum, 89+; pyxidata, 89+; rangiferina, 340+; robbinsii, new to Texas, 532; stellaris, 340+; subapodocarpa, 330; uncinalis, 340; verticillata var. crevicornis, 92
Clasmatocolea notophylla, 220+; **verrucosa**, 220+, 221*, 222*
Clasmatodon parvulus, 164+, 203
Clathroporina amygdalina, 12
cleavage patterns, in sporogenesis, 141, 159
Climacium americanum, 164+, 204, 444; dendroides, 73, 444
CLINTON, G. W., 128
Coccocarpia cronia, 30+
Coccomyxa, 246
coevolution, in lichen-herbivore, 332
Coleochaetium scaberrimum, 419; standleyi, 419
Collema conglomeratum, 30; subflaccidum, 30+
Cololejeunea biddlecomniae, 164+; bokorensis, 278; macounii, ecology, 227
Colorado, Anacolia menziesii in, 253
Committee to review goals, purpose, and operations of ABLS, 129
compensation levels, in lichens, 505
competitive interactions, in cryptogams, 333
Conocephalum conicum, 96, 353
Conostomum tetragonum, 359
constictic acid, 331+
Cornicularia aculeata, 92; divergens, 92; loxensis, 529; muricata, 92
cortical tissue, in lichens, 332
corticolous lichens, 281, 296
Costesia, 467+; character states in, 472+; phytogeography, 471; spongiosa, 467, 474*
Coumassie brilliant blue, 340+
cover, bryophytes in sinks, 162+
CRANDALL-STOTLER, B. *See* STOTLER, R. E. & B. CRANDALL-STOTLER, 111, 258, 370, 592
CRANDALL-STOTLER, B. & W. R. GUERKE. Developmental anatomy of Jubula Dum. (Hepaticae), 179
Cratoneuron commutatum, 204
CROSBY, M. R., Recent literature on mosses—114, 118; —115, 270; —116, 388
CROSBY, M. R. *See* HERNSTADT, I. C., C. HEYN & M. R. CROSBY, 536
CROSBY, M. R. & C. R. BAUER. Recent literature on mosses—117, 603
Crossidium crassinerve, capsules as food source for ants, 524; squamiferum var. pottioideum, 257
Cryphaea, 455; brevipila, 452*; glomerata, 204, 452*; ravenelii, 204
Cryphaeaceae, 451+
Cryphidium, 455; leucocoleum, 452*
Cryptocarpon, 405+
Crypto-forsstroemia, 455+
cryptogam, competitive interactions in, 333
Cryptoleptodon, 455; flexuosus, 454*, 458*
β-cryptoxanthin, 22+
Ctnedium molluscum, 204
CULBERSON, W. L. (REVIEW), H. A. BELTMAN, Vegetative struckturen der Parmeliaceae und ihre Entwicklung, 402
Cyclodictyon albicaule, 556; lindgianum, 556; varians, 556
Cynodontium tenellum, 238
Cyrtidula americana, 19
Cyrtomnium hymenophylloides, 238
cytokinin, exogenous effect on Riccia, 67
cytomictic channels, 145
cytoplasmic lobing, in sporogenesis, 141
CZECZUGA, B. Investigations on carotenoids in Embryophyta. I. Bryophyta, 21

Dactylina arctica, 88+; ramulosa, 88+
Daltonia longifolia, 556
Dawsonia longiset, 543*; polytrichoides, 542*, 543*; superba, 73, 542+, 543*
DE BONNEVAL, L., 87
DELGADILLO M., C. *See* ZANDER, R. H., C. DELGADILLO M. & P. M. ECKEL, 508
Dendroalsia, 455
Dermatocarpon miniatum, 340+
Desmatodon ellesmerensis, 489; latifolius, 359, var. muticus, 361; laurei, 361; leucostoma, 254; obtusifolius, 204, 365; plinthobius, 204, 356; systylius, 359
developmental anatomy, in Jubula, 179+
Dichelyma falcatum, 238
Dichodontium pellucidum, 238
Dicnemonaceae, 474
Dicranaceae, 526, 559
Dicranella, 559; grevilleana, 361; heteromalla, 204, sensitivity to trampling, 301; rufescens, 204; varia, 204, 492
Dicranum, 280, sensitivity to trampling, 302; subg. Trichodontium, 589; angistirete, 582; aucklandicum, 589+, 590*; donnellii, 582; flagellare, propagules, 483; fulvum, 307, 330; fuscescens, 204; montanum, 307; **pseudofalcatum**, 590+, 591*; sabuletorum, 204; scoparium, 204, 306, chlorophyll a/b ratio, 85; subleucogaster, 581+; zygodonticarpum, 582
Didymodon sect. Vineales, 230; acutus, 230, var. acutus, 493; brachyphyllus, 256; fallax var. fallax, 230, var. recurvifolia, 230 var. **reflex-**

us, 230; jamesonii, 367; rigidicaulis, 230; rigidulus, 230; tophaceus, 204
diffuse stage, in moss sporogenesis, 143*, 146+
Diphyscium foliosum, 204, 238, 306, 372, as food source for ants, 525, new to Mississippi, 372
Diplophyllum, 497, distribution in Western North America, 497; albicans, 96, 497+; andrewsii, 497+; apiculatum, 307, 497+, 503; hyalinum, 503; imbricatum, 497+, North American distribution, 500*; microdontium, 497+, North American distribution, 501*; obtusatum, 497+; obtusifolium, 97, 487+; plicatum, 497+, North American distribution, 502*; subg. Austrodiplophyllum, 497+; subg. Diplophyllum, 497+; subg. Macrodiplophyllum, 497+; subg. Protodiplophyllum, 497+; taxifolium var. macrostichum, 498+, North American distribution, 499*, var. taxifolium, 56, 97, 498+
diplotene, 148
Directory of bryologists and bryological research, new edition, 129
dispersal, by birds, 493
Distichium capillaceum, 493; hagenii, 359; inclinatum, 359
Ditrichaceae, 526
Ditrichum aucklandicum, 589; brevirostre, 589; pallidum, 204, 545+, archesporial cell in, 141, meiotic prophase, 137+, regeneration in, 309, sporogenesis, 137+, 153+; pusillum, 306; rufescens, 556
Donrichardsia macroneuron, 512+, 514*, culturing, 517, phenetic affinities, 512+
Douinia, 497; ovata, 97, 227
Drepanocladus aduncus, 204, var. capillifolius, 492; exannulatus, 492+; lycopodioides, 492; sendtneri, 493; trichophyllus, 492+; tundrae, 492; vernicosus, 492
Driftless Area refugium, 535
Drummondia, 441, 466; prorepens, 204

ECKEL, P. M. *See* ZANDER, R. H., C. DELGADILLO M. & P. M. ECKEL, 508
ECKEL, P. M. *See* ZANDER, R. H. & P. M. ECKEL, 209
editorial, 617
Edrudia constipans, 64+, 65*
Edwards Plateau, Texas, 512+, historical geography, 519
effect of coal-fired generating station on corticolous lichens, 281
EGAN, R. S., 279
EGAN, R. S. Recent literature on lichens—104, 103; —105, 263; —106, 380; —107, 596
EGAN, R. S. (review), H. KROG, H. ØSTHAGEN & T. TØNSBERG. Lav Flora. Norske bruskog bladlav, 614
EGAN, R. S. (review), J. W. THOMSON. Lichens of the Alaskan Arctic Slope, 612
Encapypta brevicolla, 356; longicolla, new to Newfoundland, 563; rhaptocarpa, 444; streptocarpa, 238; vulgaris, 444, var. mutica, 209
energy investment, in asexual reproduction, 344
ENGEL, J. J. Austral hepaticae. XII. A new species of Clasmatocolea (Hepaticae) from Tasmania, 220.
Entodon cladorrhizans, 164+, 204; concinnus, 238; macropodus, 204; seductrix, 204
Entosthodon leibergii, 356; tucsonii, new to Colorado, 257
Eotrichocolea polyacantha, 81
Ephemerum crassinervium, 356; serratum, 356
Erpodiaceae, 474
Erpodium domingense, 556; glaziovii, 556
Erpyrenula leucoplaca, 9
Erythrobarbula binnsii, 367
Erythrodontium tenuicaule, 459*, 460
ESSLINGER, T. L. Typification of Oropogon loxensis and description of two related species, 529
Eucladium verticillatum, 204, 517
Eurhynchium, 512; hians, 164, 204, 513; macroneuron, 512; pulchellum, 204, 492, var. barnesii, 513, var. pulchellum, 513; riparioides, 512+
Eurohypnum leptothallum, 461
European Trichostomeae, 132
Evernia mesomorpha, 288
evolution of algal-fungal relationship in appressorum-type, 177
excurrentinervis-type peristome, 323
exine, in moss spores, 550
2,4-D, exogenous effect on Riccia, 67

Fabronia ciliaris, 204; gymnostoma, 204
FAHSELT, D. Alternative method for analyzing protein in lichens, 340
field microscope light, 404
FIFE, A. J. The affinities of Costesia and Neosharpiella and notes on the Gigaspermaceae (Musci), 466
FIFE, A. J. & G. R. BRASSARD. New or additional moss records from Newfoundland V, 358
Fissidens, 559+; adiantoides, 204, 238, 370, peristome, 315+, 321*; asperifolius var. cambodiana, 278; asplenioides, 556, peristome, 315+, 322*; bryoides, peristome, 315+, 320*; bushii, peristome, 315+, 321*; cristatus, 164+, 204, 556; diplodus, 256, var. **richardsii**, 256; elegans, 559; excurrentinervis, peristome, 315+, 320*; exiguus, 164; exilis, peristome, 315+, 318*; flexinervis, 528; fontanus, 204; gladiolus, 528, 560+, 561*, peristome, 315+, 319*; guianensis, 556; intermedius, 526; limbatus, 145, 553, peristome, 315; manateensis, peristome, 315+, 322*; milobakeri, peristome, 315+; mollis, 556; muriculatus, 256; obtusifolius, 204, peristome, 315+, 319*; osmundoides, 204;

pauperculus, 528; polypodioides, 556, peristome, 315+, 322*; prionodes, 528, peristome, 315+, 318*, 319*; radicans, peristome, 315+, 318*; scariosus, 559, peristome, 315+, 318*; subbasalaris, 164+, 204; subulatus, 526, 527*, 559, in Brazil, 526, peristome, 315+, 318*, 319*, 320*; taxifolius, 164, 204, peristome, 315+, 321*; viridulus, 164+, 204; SEM study of peristome, 314; subg. Fissidens sect. Aloma, 314+, sect. Amblyothallia, 314+, sect. Bryoidium, 314+, sect. Crenularia, 314+, sect. Heterocaulon, 314+, sect. Semilimbidium, 314+, sect. Serridium, 314+, sect. Weberiopsis, 314+; subg. Octodiceras, 315+
Fissidentaceae, 559+, 565
flavonoids, in Pottiaceae, 228+
Florida, pyrenocarpous lichens, 1
Fontinaliaceae, 477
Fontinalis, 280, 521, dispersal success, 478, interspecific competition, 478, rhizoid production, 477, stoloniferous branches, 479+; antipyretica, 478+, carotenoids, 24+; dalecarlica, 478+, stoloniferous branches, 482*; duriaei, 478+; filiformis, 205; gigantea, 238; hypnoides, 478+; neomexicana, 478+; novae-angliae, 205, 477+, var. cymbifolia, 369, var. novae-angliae, 369; squamosa, carotenoids, 22, 24, 25; sullivantii, 205, 369
Forsstroemia, 451+; sect. Microforsstroemia, 457; sect. Cryptoforsstroemia, 457; australis, 457; cordata, 457; coronata, 453+; cryphaeoides, 457; cryphaeopsis, 457; cuspidata, 457; dixonii, 457; schensiana, 456; felipponei, 457; flagellacea, 457; inclusa, 457; incrassata, 457; indica, 455+, 456*; japonica, 457; julacea, 457; kusnezovii, 457; lasioides, 457; **leptodontoidea**, 451+, 452*; mandschurica, 457; neckeroides, 457; neomexicana, 457; nitida, 457; occulata, 457; ohioensis, 457; paraguensis, 457; producta, 457; recurvimarginata, 457; rigida, 451+, 452*; secunda, 457; sinensis, 451; stricta, 457; subcoronata, 457; subproducta, 457; **thomsonii**, 456*, 457; trichomitria, 205, 453+; tripinnatum, 457; ulei, 453+
FRAHM, J. Synopsis of the genus Campylopus in North America north of Mexico, 570
FREDSKILD, B., 87
Frullania, 84; brittoniae, 164+; riparia, 164+; tamarisci subsp. nisquallensis, 97, 227; squarrosa, 164
frullanioid characters, in Jubula, 179
FULFORD, M. Awarded Certificate of Merit, 129
Fulgensia, 66
fumarprotocetaric acid, 341
Funaria americana, 205, 335+, distribution, 337+, geography, 338; flavicans, 205; hygrometrica, 238, 311, 545, carotenoids, 24+; mediterranea, 337; serrata, 205
Funariaceae, 467+
Garysmithia bifurcata, 458*, 461
Geocalyx graveolens, 97
Gigaspermaceae, 467+, 540, character states in, 472+, comparison of genera, 470, phygeography, 471
Gigaspermum, character states in, 472+, distribution, 540+, in Israel, 536, phytogeography, 471, 540; mouretii, 471, 536+, 538*; repens, 469*, 475+, 536+, 538*; subrepens, 471, 537+; tumidum, 536+
GLIME, J. M. Effects of temperature and flow on rhizoid production in Fontinalis, 477
Glossadelphus, 463
glucans, 340
glyrophoric acid, 331
GODFREY, G. A. *See* GODFREY, J. D. & G. A. GODFREY, 224
GODFREY, J. D. & G. A. GODFREY. Notes on hepatics from the Pacific Northwest, 224
Gondwanaland distribution type, 474, flora, 445
Graphis scripta, 288+
grazing activity by Pallifera varia, 328
GREEN, T. G. A. *See* SNELGAR, W. P. & T. G. A. GREEN, 505
Greenland, lichens, 87
Grimmia alpestris var. holzingeri, 356; alpicola, 205; anodon, 365; apocarpa, 164+, 205, 320, var. conferta, 205; arizonae, 205; atricha, 365; elongata, 253; hartmanii var. anomala, 361; laevigata, 205, chlorophyll a/b ratio, 85+; occidentalis, 253; olneyi, 205; pilifera, 205; pulvinata, carotenoids, 24, 25; rauii, 205; torquata, 238; wrightii, 205
Groutiella, 408
GUERKE, W. R. *See* CRANDALL-STOTLER, B. & W. R. GUERKE, 179
Gymnocolea inflata var. heterostipa, 97, var. inflata, 97
Gymnomitrion concinnatum, 238; corallioides, 351+; obtusum, 97
Gymnostomum, 280; aeruginosum, 163+, 205, 489; calcareum, 205; recurvirostrum, 238
Gyrothyra underwoodiana, 97

Haematomma lapponicum, 92; puniceum var. subinnatum, 219
HALE, M. E., JR. The lichen genus Relicina (Parmeliaceae) in India and Sri Lanka, 77
HANSEN, E. S. Lichens from northwestern Greenland collected on botanical expeditions in 1975 and 1977, 87
Haplocladium microphyllum, 205, 556; virginianum, 306
Haplohymenium triste, 164+, 205
Haplomitrium hookeri, 97
Harpanthus scutatus, 225
HARPEL, J. D. Orthotrichum pumilum new to California, 201

HARRIS, R. C. *See* S. C. TUCKER & R. C. HARRIS, 1
harvester ant, 524
HATTAWAY, R. A. The calciphilous bryophytes of three limestone sinks in eastern Tennessee, 161
HEDWIG, J., 335
Hedwigia ciliata, 205, 330
Helicodontium capillare, 556
hepatics, carotenoids, 22; of North Cascades Range, 94; of Pacific Northwest, 224; of Tasmania, 220; of Washington, 94; recent literature, 111, 263, 373, 592
Herberta aspara, 278
Herbertus aduncus subsp. aduncus, 97
herbivore, polyphagous, 333
HERMAN, F. J. Distinctions between Anacolia menziesii (new to Colorado) and Bartramia stricta, 253
HERNSTADT, I. *See* LORIA, M. & I. HERNSTADT, 524
HERNSTADT, I., C. C. HEYN & M. R. CROSBY. New data on the moss genus Gigaspermum, 536
Herpetineuron toccoae, 205
Heterodermia tremulans, 288
HEYN, C. C. *See* HERNSTADT, I., C. C. HEYN & M. R. CROSBY, 536
Hispaniola, moss flora, 555
historical geography, Edwards Plateau, 519
HOE, W. J. *See* VITT, D. H. & W. J. HOE, 212
Holodontium falcatum, 589
HOLZINGER, J., 335
Homalia trichomanoides, 164+, 238
Homalotheciella subcapillata, 164
Homalothecium leskeoides, 556; sericeum, 238
Homomallium adnatum, 164+, 205, 307
HONG, W. S., A study of the distribution of Diplophyllum in Western North America, 497
HONG, W. S. Hepaticae of the North Cascades Range, Washington, 94
HONG, W. S. Hepaticae of Rocky Mountain National Park, Colorado, 351
HONG, W. S., The genus *Scapania* in western North America. II. Taxonomic treatment, 40
Hookeria lucens, 521
HOPKINS, A. W. *See* JACKSON, D. W. & A. W. HOPKINS, 532
Horridohypnum, 463
Horridohypnum mexicanum, 462*, 463
Huilia albocaerulescens, 330+
Husnotiella revoluta, 205, var. palmeri, 205
Hygroamblystegium, 512; fluviatile, 480, var. fluviatile, 513, var. ovatum, 513; noterophilum, 205, 513; tenax, 164+, 205, var. spinifolium, 205, 513, var. tenax, 513
Hygrohypnum eugyrium, 359; subeugyrium, 359; montanum, 359; smithii, 254, 359
Hylocomnium brevirostre, 238; splendens, 238, carotenoids, 24+, sensitivity to trampling, 302, var. tenue, 461
Hymenostomum breutelii, 556; tortile, 536
Hymenostylium recurvirostrum, 229
Hyophila involuta, 164, 205
Hypnobryales, 460
Hypnodendraceae, 523
Hypnum, 280; bambergii, fossils, 486; circinale, diffuse stage in sporogenesis in, 149; cupressiforme, 205, carotenoids, 24+; curvifolium, 164, 205; imponens, 205, 308, sensitivity to trampling, 301; pallescens, 205, 306; rusciforme, sporogenesis in, 159; polypterum, 556; vaucheri fo. tereticaulis, 461
Hypogymnia austerodes, 92; oroarctica, 92; subphysodes, 83
Hypotrachyna osorioi, 219

importance values, bryophytes in sinks, 162+
inhibitory chemistry, investment in, 333
intergeneric hybrids, 159
IRELAND, R. R. The taxonomic status of Ceratodon heterophyllus, 234
isoelectricfocussing, 340
Isoetes, amoeboid plastid in, 147
isolichenin, 340
Isopterygium elegans, 205, 307; subtenerrimum, 557; tenerum, 205
Isothecium myosuroides, 238; stoloniferum, 243
Israel, moss flora, 279

JACKSON, D. W. & A. W. HOPKINS. New additions to the lichen flora of Texas and the United States, 532
Jamesoniella autumnalis var. autumnalis, 97
JANSSENS, J. A. & R. H. ZANDER. Leptodontium flexifolium and Pseudocrossidium revolutum as 60,000 year-old subfossils from the Yukon Territory, Canada, 486
JORDAN, W. P. **Edrudia**, a new genus from California (Lichenes: Teloschistaceae), 64
Jubula, antheridial development in, 194, archegonial development in, 195, developmental anatomy in, 179+, frullanioid characters in, 179, leaf formation in, 187+, lejeuneioid characters in, 179, protonema in, 197, Radula-type subfloral innovations in, 179, spore germination in, 197+, sporophyte anatomy in, 195+, sporophyte development in, 195, stem formation in, 191+, systematic position of, 198+; blepharophylla, 182+, 182*, 184*; japonica, 180+, 180*, 182*, 184*, 186*, 190*, 192*; pennsylvanica ssp. bogotensis, 185+, 188*, 190*, ssp. hutchinsiae, 180+, 180*, 182*, 183*, 184*, 186*, 188*, 190*, 192*, ssp. javanica, 180+, 180*, 182*, 183*, 190*, ssp. pennsylvanica, 180+, 180*, 182*, 183*, 184*, 186*, 188*, 190*, 192*

JUDD, W. S. Notes on the moss flora of Hispaniola, 555
Julocladium, 467
Jungermannia atrovirens, 97; exsertifolia, 97, subsp. cordifolia var. cordifolia, 352; gracillima, 97; hyalina, 97, 307; leiantha, 97; mollissima, 79+; obovata, 97; pumila, 97; rubra, 97

KEENER, C. A. (review), R. J. LITTLE & C. E. JONES, A dictionary of botany, 615
ketocarotenoid, 22
Kiaeria blytii, 361; starkei, 254
kinetin, exogenous effect on Riccia, 67
Kurzia sylvatica, 307
KUWAHARA, Y., 279

Labrador, bryophyta exsiccata from, 237
LAMBRIGHT, D. D. & S. C. TUCKER. Observations on the ultrastructure of Trypethelium eluteriae Spreg., 170
lamellae, 315+
lampbrush chromosome, 147
LANE, D. N. *See* WYATT, R., D. M. LANE, & A. STONEBURNER, 247
LANGLOIS, A. B., 1
LARZARENKO, A. S., death, 278
Lasallia papulosa, 330+
Lasia orthotrichoides, 410*
Laurasian flora, 445
Laurera megasperma, 7
LAWREY, J. D. Correlations between lichen secondary chemistry and grazing activity by Pallifera varia, 328
LAWREY, J. D. Sexual and asexual reproductive patterns in Parmotrema (Parmeliaceae) that correlate with latitude, 344
leaf formation, in Jubula, 187+
leaves, regeneration from, 542
Lecanora atrosanguinea, 71; bolanderi, 66; campestris, 330; conizaea, 288; constipans, 64; epibryon, 89+; hypoptoides, 288; melanophthalma, 90+; novomexicana, new to Texas, 533; polytropa, 92; varia, 288
Lecidea atrobrunnea, 90+; lapicida, 92
Leiomitrium scaberrimum, 419
lejeuneioid character, in Jubula, 179
LEMMON, B. E. *See* BROWN, R. C. & B. E. LEMMON, 137, 153, 545
Lepicolea ochroleuca, 84
Lepidopilum mueller, 577; scabrisetum, 557
Lepidozia, 84; reptans, 97, 353
Lepista nuda, 330
Lepraria membranacea, 288+, 535; neglecta, 92, 288; zonata, 330
Leprocaulon subalbicans, 89+
Leptangium, 536, imbricatum, 475
Leptobryum pyriforme, 205, 492
Leptodon, 453+
Leptodontaceae, 453+; subfam. **Forsstroemioideae**, 453
Leptodontoideae, 455
Leptodictyum riparium, 205, 493; trichopodium, 205, 492
Leptodontium flexifolium, 230, 490+, dispersal history, 493, distribution, 494*, propagula, 493, subfossils, 486+; viticulosoides, 230, var. panamense, 557
Leptopterigynandrum austro-alpinum, 458*, 459*, 460; **clavatum**, 459*, 460+; decolor, 459*, 460; strictcaule, 461; subintegrum, 461
Leptogium chloromelum, 30+; corticola, 30; cyanescens, 30+
Leptostomaceae, 474
Leptotheca, 475
Leptotrema **desquamans**, 368; **oligospermum**, 368
Lepyrodon tomentosus, 557
Lepyrodontopsis trichophylla, 557
Leskea, 280; australis, 205; gracilescens, 164, 205; obscura, 206; polycarpa, 206
Leskeella arizonae, 230
LESQUEREUX, L., 128, 335
Leucobryum albidum, 206, 370; antillarum, 557; crispum, 557; glaucum, 206, 238, carotenoids, 24+, chlorophyll a/b ratio, 85, sensitivity to trampling, 301; incurvifolium, 370, in Florida, 370; polakowskyi, 557
Leucodon brachypus, 206, var. andrewsianus, 361; julaceus, 164, 206, chlorophyll a/b ratio, 85; thomsonii, 455+
Leucodontaceae, 451+, 455; subfam. Forsstroemioideae, 453
Leucodontopsis, 453
Leucolepis menziesii, 521
Leucoloma crugerianum, 557; serrulatum, 557; tortellum, 250
Leurera madreporiformis, 9
LEWIS, M. Field microscope light, 404
lichen communities, as bioindicators, 281, 296
lichen-herbivore, coevolution in, 332
lichenin, 340
lichens, analyzing proteins in, 340; carbon dioxide exchange in, 505; carotenoids, 27; corticolous communities, 82; gray-beech forest, 36; in Greenland, 87; in North Carolina, 29; K-selection in, 349; morphological changes in, 296+; nitrogen fixing, 29; on bones, 90; on fell-fields, 90; on fens, 90; on snowbeds, 90; photorespiration in, 505; recent literature, 103, 263, 380, 596; r-selection, 349; reproductive strategy in, 347; saxicolous, 90, 329; secondary chemistry, 328; SO_2 effects on morphology, 296+; SO_2 sensitive, 281+; southern Appalachians, 29; southern Chile, 82; stress adaptation, 349; surface area, 32+; Valdivian forests, 82; from Argentina, 219; from Florida, 1; from Louisiana, 1
Limbella tricosta, 521+
limestone sinks, in Tennessee, 161+
limestones, Silurian, 338
Lindbergia brachyptera, 206
line-intercept transect, 162

lipid bodies in sporocyte, 138+
Lithothelium, 10
Lobaria scrobiculata, 246; oregana, 32, 37; pulmonaria, 30+; quercizans, 30+
LODWICK, L. D. & J. A. SNIDER. The distribution of Sphagnum in Texas, 214
loesslands, Wisconsin, origin of, 337*
Lopadium pezizoideum, 89+
Lophocolea bidentata, 98, 164+; cuspidata, 164+, var. cuspidata, 98; heterophylla, 98, 164, 307, 521, 553; minor, 98; ventricosa var. ventricosa, 98
Lophozia alpestris, 98, 351+; ascendens, 98, 351+; collaris, 98, 353; gillmanii, 98; guttulata, 98, 353; heterocolpos, 98; incisa, 351+, var. incisa, 98; longidens, 98, subsp. longidens, 353; menzelii, 98; obtusa, 353; opacifolia, 98, 353; ventricosa, 351+, var. longifolia, 98, var. silvicola, 98, 353
Lorentiziella, 470, character states in, 472+; glauca, 475; globiceps, 476; imbricata, 469*, 475+, 540; paraguensis, 476
LORIA, M. & I. HERNSTADT. Moss capsules as food of the harvester ant, Messor, 524
Louisiana, Pyrenocarpous lichens, 1
lutein, 22+, epioxide, 22+
lycopene, 22+

Macrocoma, taxonomy, 405, typification of names, 405; subg. Macrocoma, 407+, world distribution, 446*; subg. Trachyphyllum, 407+; **abyssinica**, 408, 419+, 422*, 427+, 447, world distribution, 446*; brasiliensis, 408, 409, 447, world distribution, 446*; capillicaule, 409; filiforme, 407+, 418; frigida, 408+, world distribution, 446*; gastonyi, 408+, 411*, 447, world distribution, 446*; **gracillima**, 433, 434*; hyalinum, 419+, 427+; intricata, 417; **iwatsukii**, 408, 422+, 430*, world distribution, 446*; lycopodioides, 408, 423+, 424*, world distribution, 446*; orthotrichoides, 408+, 412*, world distribution, 446*; **papillosa**, 433, 434*; protractum, 421, pulchella 408, 419*, 425+, world distribution, 446*; scaberrima, 419; sullivantii, 437+, geographical variation in, 437+; tenue, 409+, subsp. **sullivantii**, 408, 413, 416*, 425+, leaf variation in, 440*, world distribution, 446*, subsp. tenue, 408, 424*, 425+, leaf variation in, 440*, world distribution, 446*; virescens, 421
macrofossils, 496
Macromitrio dregeri, 417
Macromitrium, 405+; subg. Trachyphyllum, 433; abyssinicum, 419, 420*; anacamptophyllum, 418; angulicaule, 419, var. phyllorhizans, 413; bolanderi, 420*; bolivianum, 418; brasiliense, 409, 410*; braunioides, 418; caespitans, 428; capillicaule, 410*; chrysomitrium, 410*, 413; cirrosum, 557; confusum, 426*, 428+; consanguineum, 420*, 432; contractum, 432; dawsoniaemitrium, 420*, 429+; dregei, 415+; eucalyptorum, 426*, 428, var. recurvulum, 429; fitzgeraldii, 418; frigidum, 409, 410*; gastonyi, 417; geheebii, 426*, 428+; ghiesbreghtii, 417, var. brevifolium, 418; gracillimum, 433; hyalinum, 421, 426*, 429; hymenostomum, 417; intricatum, 414*, 417+; johnsonii, 428; krausei, 418; lampromitrium, 418, 420*; lieboldtii, 414*, 417; liliputanum, 432; lycopodioides, 419+; macropyxis, 410*, 423; mexicanum, 414*, 417; novaevalesiae, 426*, 429; okamurae, 420*, 432; orthotrichoides, 419; papillosum, 433; paraphysatum, 414*, 417, var. chilense, 414*, 418; perottetii, 414*, 431+; phyllorhizans, 410+; pringlei, 410*; progressum, 414*, 417; protractum, 420*, 421+; pulchellum, 424; pycnangium, 418, 420*; quinquefarium, 410*, 419; recurvulum, 429; sobrinum, 426+; squarrulosum, 414*, 432+; subpycnangium, 410+; sullivantii, 413+, 414*, 439+; tenue, 415+, fo. microphyllum, 427, subsp. dregei, 428, subsp. sullivantii, 409+, 416*, var. brachypus, 428, var. lycopodioides, 423; virescens, 420*, 421
MAHLER, B. D. & W. F. MAHLER. Checklist of mosses of Oklahoma, 202
MAHLER, W. F. 278
MAHLER, W. F. *See* MAHLER, B. D. & W. F. MAHLER, 202
MAKHIJA, U. *See* PATWARDHAN, P. G. & U. MAKHIJA, 368
manateensis-type peristome, 323
marbles, Silurian, 338
Marchantia, 150; polymorpha, 98, 353, carotenoids, 21, 23
Marsupella brevissima, 98; emarginata, 238, subsp. emarginata var. emarginata, 98; sphacelata fo. media, 99, 353; sparsifolia, 99
MARTIN, C. E. Chlorophyll a/b ratios of eleven North Carolina mosses, 84
Maschalocarpus ecklonii, 426+
MAURICE LOUIS JEAN BIZOT (1905–1979), 564, portrait, 565, publications, 565+
Meesia, 280; uliginosa, 238
Meiotic spindle, 141
Melanotheca achariana, 9; aggregata, 9; anomala, 9, 175; concatervans, 15; cruenta, 9, 175
Melophyllum radiculosum, 255
Menegazzia albida, 83; dispora, 83
Meteorium illecebrum, 557; remotifolium, 557
Metzgeria conjugata, 99, 521; furcata, 238; temperata, 227
Mexican Highlands, 519
Mickel cell homogenizer, 342+
Miconia plukenetii, 256
Microforsstroemia, 455+
microhabitats, in sinks, 165
microtubules in sporogenesis, 148*, 150+, 154*
Mielichhofera macrocarpa, 366; mielichhoferii, 365; nitida var. gymnostoma, 366; schiedeana, 366; **tehamensis**, 355+, 366*
Mississippi, Diphyscium foliosum in, 372

mitochondria, in sporogenesis, 143, 155
Mittenothamnium, 461; diminutivum, 557; **sharpii**, 461
Mnium affine, 206, carotenoids, 22+; ciliare, 206, gamete dispersal, 73+; cuspidatum, 75, 164, 206, 553, carotenoids, 24+, chlorophyll a/b ratio, 85; hornum, 141+, 307; marginatum, 164+; medium, 164+, 206; punctatum, 206; rostratum, 164+, 206; rugicum, 206; undulatum, carotenoids, 24+
MOBERG, R. Anaptychia ulotrichoides new to North America, 251
Moenkeymeyera richardsii, 256
moisture levels, in sinks, 165
Molendoa sendtneriana, 492
Monoblastia lutescens, 10; palmicola, 10; rappii, 10
Morinia, key to taxa, 508; crassicuspis, 508; ehrenbergiana var. ehrenbergiana, 510+, var. elongata, 510; **saitoana**, 508+, 509*
moss capsules, as food for ants, 524
moss sporogenesis, diffuse stage in, 143*, 146+
mosses, chlorophyll a/b ratios, 84; North Carolina, 84; recent literature, 118, 270, 388, 603; septal formation in, 159; spore separation in, 159; of Oklahoma, 202+
MSE Sonicator, 342
Mt. baker-Snoqualine Natl. Forests, hepaticae of, 95
MUELLER, D. M. J. (review), Advances in botanical research, Vol. 6, 616
Muelleriella, 444
MUHLENBERG, H. L. 335
Musci, carotenoids, 22
mutatochrome, 22+
mutatoxanthin, 22+
Mycoporellum eschweileri, 19; lacteum, 19
Mylia anomala, 99; taylori, 99
Myurella julacea, 238, 361; sibirica, 164+, 238; tenerrima, 238

Nanobryaceae, 559+
Nanobryum duemmeri, 559+; gladiolum, 559
Nardia compressa, 99; geoscyphus var. geoscyphus, 99; insecta, 225; japonica, 224+, ecology, 225; scalaris var. scalaris, 99
Neckera, 280; complanata, 238; domingensis, 455; douglasii, 521
Neckeraceae, 455, 523
Neomacounia, 457; nitida, 458*
Neosharpiella, 467+, character states in, 472+, phytogeography, 471; aztecorum, 467+, distribution, 474*; turgida, 468+, 469*, distribution, 474*
neoxanthin, 22+
nephrin, 243+
nephroarctin, 243+
Nephroma antarcticum, 83, 246; australe, 246; cellulosum, 246; gyelnikii, 246; helveticum, 30+; lepidophyllum, 246, f. hypomelaena, 246; microphyllum, 246; **occultum**, 243+, 244*, 245*; parile, 30+
Neptune State Park, Oregon, mosses of, 355
Neskowin Crest Research Natural Area, Oregon, mosses of, 355
NEUMANN, A. J. & A. F. ROGERS. Contribution to the bryoflora of Texas, 369
neurospcrene, 26
Newfoundland, bryophyte exsiccata from, 237
Nipponolejeunea, 179
NOBLE, W. J. Catillaria **columbiana**, comb. nov., from western North America, 71
norstictic acid, 331, 342
North America, Anaptychia ulotrichoides in, 251
North Carolina, mosses of, 84
North Cascades National Park, Hepaticae of, 94
Nostoc, 243+
Nowellia curvifolia, 307
nuclear pores, 142*
nucleolus in meiosis, 143*, 144*, 146*
numerical analysis, 513+; taxonomy, in mosses, 518

obtusifolius-type peristome, 323
Ochrolechia yasudae, 330
Octoblepharum erectifolium, 557
Odontoschisma denudatum, carotenoids, 23; sphagni, 238
Oedipodiella, 470; character states in, 472+
Oedipodium australe, 476
oil bodies, in sporocyte, 141
Okanogan Natl. Forest, Hepaticae of, 95
Oklahoma, mosses of, 202+
Oligotrichum hercynicum, 361
Omphalodiscus decussatus, 90+
Oncophorus virens, 493
Opegrapha pulicaris, 288+
Oregon, moss flora of, 355
Oropogon **atranorinus**, 530; formosanus, 531; **lopezii**, 531; loxensis, typification, 529, var. atroalbicans, 530
Orphniospora atrata, 92
Orthotrichum anomalum, 444; cupulatum, 206, 445; diaphanum, 206; flowersii, 201; gymnostomum, 238; hornschuchii, 424; lyellii, 201; microphyllum, 426*; ohioense, 206; pumilum, 206, in California, 201; pusillum, 206; pylaisii, 238; stellatum, 206; strangulatum, 164+, 206; strictum, 254; tenellum, 201, 425+, 426*, 439
osmiophilic granules, in plastids, 551
OSORIO, H. S. Contribution to the lichen flora of Argentina XII. New or additional records from Buenos Aires Province, 219
Ozark Plateau, 519

pachynema, 143
pachytene, 143+
Pacific Northwest, hepatics of, 224
Palinocraspis, 577
Pallifera varia, food preferences, 328

Pannaria rubiginosa, 30, 37
Palmocladium leskeoides, 206
Paludella squarrosa, 238, 433
Pangaea, 445
Papillaria nigrescens, 557
Paraleucobryum, 581
Parathelium cuyabense, 10, 11*; subferruginеum, 10, 11*
Parmelia subg. Amphigymnia, 345; aurulenta, 288; bolliana, 288+, 296+, SO_2 effects on, 296+; borrerioides, 219; caperata, 288+, 296+, SO_2 effects on, 296+; centrifuga, 92; cumberlandia, 340+; disjuncta, 92; flaventior, 288; galbina, 288; hypoleucites, 219; infumata, 90+; microsticta, 219; omphalodes, 92, 330; pseudosulcata, 83; reticulata, 83; rudecta, 288+, 330; saxatalis, 92, 288+; squarrosa, 288; subaurifera, 288; sulcata, 90+, 288+; ulophyllodes, 288
Parmeliella corallinoides, 30+; manipurensis, 533, 534*
Parmelina simplicior, 534; wallichiana, 534
Parmotrema, apothecia in, 345, asexual reproductive patterns in, 344, sexual reproductive patterns in, 344; crinitum, 330; mordenii, new to Texas, 533, new to the United States, 532; uruguense, 220
Pasayten Wilderness Areas, Hepaticae of, 95
PATWARDHAM, P. G. & U. MAKHIJA. Nomenclatural note on three species of Anthracothecium, 368
PECK, J. H. Bryoxiphium norvegicum in Iowa, 535
Pellia endiviifolia, 99, 351+; epiphylla, 238; fabbroniana, carotenoids, 23; neesiana, 99, 521
Peltigera aphthosa, 92; canina, 30+, 340+; horizontalis, 30+, CO_2 compensation level in, 506; polydactyla, 30+; rufescens, 90+; scabrosa, 90+
perine, in moss spores, 551
peristome, bryoides-type, 316; excurrentinervis-type, 323; Fissidens, 315+; manateensis-type, 323; obtusifolius-type, 323; scariosus-type, 315; subbasilaris-type, 323; taxifolius-type, 323
perlatolic acid, 342
Pertusaria, 330; dactylina, 89+; panyrga, 89+; pustulata, 288+
PETERSON, W. L. & A. J. SHARP. Leucobryum incurvifolium in the United States, 370
Phascum, 280; cuspidatum, 206
phenarctin, 243+
Philonotis angulata, 557; fontanus, 206; longiseta, 206; marchica, 206; uncinata, 557
Phlyctis speirea, 71
photorespiration, in lichens, 505
photosynthesis, C_3, 505, C_4, 505
phragmoplast, in sporogenesis, 157
Phyllogonium fulgens, 557
Physcia aipolia, 288; caesia, 91, 92; carassensis, 220; chloantha, new to Texas, 533; millegrana, 288+, 296+, SO_2 effects on, 296+; orbicularis, 288+, f. albociliata, 288+, orbicularis f. rubropulchra, 288+; rudecta, 296+, SO_2 effects on, 296+; stellaris, 288+
Physciopsis adglutinata, 288; syncolla, 288
Physcomitrium subg. Julocladium, 467+; breutelii, 476; pyriforme, 206, 545
Physconia muscigena, 89+; grisea, 288
phytoene, 26
phytofluene, 26
Pilopogon, 577; gracilis, 557
Pilotrichella flexilis, 557; hexasticha, 557
Pilotrichum flagelliferum, 453+
Placodium constipans, 64
Plagiocarpa subg. **Pyrenulopsis**, 10; hyalospora, 10; **langloisii**, 10, 11*
Plagiochila, 84; austini, 164+; porelloides, 96, 164+, 354; satoi, 99
Plagiomnium ellipticum, 481; rhynchophorum, 557
Plagiopus oederiana, 238
Plagiothecium denticulatum, 306
plastids, in moss sporogenesis, 145+, 146*, 155, in spores of Leptodictyum riparium, 545+, multiplication, 553
Platismatia glauca, 83
Platydictya confervoides, 161+; subtile, 206
Platygyrium repens, 206
Pleistocene glaciation, southern limits, 337*, sediments, 487
Pleuridium, 254; subulatum, 206
pleurocarpous mosses, effects of trampling, 311; rhizoids, 481
Pleurochaete squarrosa, 206
Pleuroclada albescens, 99
Pleurophascum grandiglobum, 536
Pleurozium schreberi, 238, carotenoids, 24+
Pogonatum brachyphyllum, 206; dentatum, 238; pensylvanicum, 206; robustum, 557; tortile, 557
Pohlia, 280, 307; atropurpurea, 206; cruda, 238; heimii, 367; lescuriana, 356; nutans, 166, 206; sphagnicola, 362
Polyblastiopsis amylospora, 7; dealbens, 7; fallaciosa, 11; sublactea, 11*, 12
Polycauliona constipans, 64; regalis, 66
Polymeridium contendens, 12; **quinqueseptatum**, 12; **subcinereum**, 12
polyphagous herbivore, 333
polysaccharides, 340, in meiosis, 141+, 147*
Polytrichum, 280, regeneration in, 309; alpestre, 444; angustatum, sensitivity to trampling, 301; commune, 206, 238, carotenoids, 24, 25, sensitivity to trampling, 302, var. perigonale, 207; formosum, carotenoids, 24+; juniperinum, carotenoids, 24+; gracile, carotenoids, 24+; juniperinum, 207, 238, 553, 557; ohioense, chlorophyll a/b ratio, 86; piliferum, carotenoids, 24+; sexangulare var. sexangulare,

357, var. vulvanicum, 356; sphaerothecium, 356
Porella bolanderi, 99; cordaeana, 99; navicularis, 99; platyphylla, 164; roellii, 99, 227
Porina acrocordioides, 10; amygdalina, 10; **heterospora**, 11*, 12; nucula, 11*, 12, var. heterospora, 12; rhaphidosperma, 13; subcinerea, 12; viridiseda, 18
Porotrichum insularum, 557
Pottia truncata, 207
Pottiaceae, 474; anthocyanins in, 228; flavonoids in, 228
Prionodon densus, 557
prolamellar body, in moss plastids, 553
Protoblastenia, 66
protocetraric acid, 331+
protonema, in Jubula, 197
protonemal growth, 310+
Pseudephebe minuscula, 92; pubescens, 92
Pseudocrossidum revolutum, 491+, distribution, 492+, 494*, subfossils, 486+
Pseudocryphaea, 453; **domingensis**, 455
Pseudocyphellaria amphisticta, CO_2 compensation level in, 506; aurata, 30; billardierii, CO_2 exchange in, 505, photosynthetic rate, 506; colensoi, CO_2 compensation level in, 506; crocata, 30+, 83; delisia, CO_2 compensation level in, 506, CO_2 exchange in, 505; dissimilis, CO_2 compensation level in, 506; hookeri, CO_2 compensation level in, 506; impressa, 83; intricata, 83; flavicans, 83; nitida, 82; psilophylla, CO_2 compensation level in, 506
Pseudocyphellaria-Nephroma lichen community, 82
Pseudocyphellaria nitida-Sphaerophorus tener-Psoroma hispidilum lichen community, 82
Pseudoleskea, 460
Pseudoleskeella arizonae, new to Colorado, 257; cantenulata, 359; tectorum, 461
Pseudoparmelia baltimorensis, 330+
Pseudopyrenula, 7; subgregaria, 13
Psora rubiformis, new to Texas, 533
Psoroma hispidilum, 82; sphinctrinum, 83
Pterigynandrum decolor, 460; filiforme, 238, var. mexicanum, 462; sharpii, 461; sinense, 451
Pterobryaceae, 455, 523
Pterogonium, 455
Pterygoneurum lamellatum, 492; subsessile, 207, 492
Pterygynium filiforme, 451+
Ptilidium californicum, 100; ciliare, 100; pulcherrimum, 100
Ptilium crista-castrensis, carotenoids, 24+, sensitivity to trampling, 302; orthothecium, 556
Ptychomitrium drummondii, 207; incurvum, 207
PURSELL, R. A. Fissidens diplodus Mitt. var. **richardsii** (R. S. Williams) Pursell, comb. nov., 256
PURSELL, R. A., 279
PURSELL, R. A. & W. D. REESE. The rediscovery of Fissidens subulatus Mitt. in Brazil, 526
PURSELL, R. A. & W. D. REESE. The taxonomic status of Nanobryaceae (Bryopsida), 559
pustulan, 340
Pylasiella selwynii, 207
Pyramidula tetragona, 207
Pyrenastrum bicolor, 13; fuscum, 13
Pyrenocarpous lichens, 1; from Florida, 1; from Louisiana, 1
Pyrenocollema **prospersella**, 13; **tichothecioides**, 13
Pyrenula **aquila**, 13, 14*; bahiana, 15; cerina, 15; **concatervans**, 15; imperfecta, 15; neglecta, 25, 288+; punctella, 15; quassiaecola, 15; **rubrostoma**, 14*, 16; santensis, 16; **texana**, 14*, 16, 17; xyloides, 16
Pyxine sorediata, 288

r-selection, in lichens, 349
Radula bolanderi, 100, 227, 521; complanata, 100; mollis, 165+
Radula-type subfloral innovations in Jubula, 179
Ramalina fastigiata, 288; intermedia, 330; sinensis, 288
RATKOWSKY, D. A. Some further synonyms of Bryoerythrophyllum jamesonii (Tayl.) Crum, 367
RATKOWSKY, D. A. Tasmanian and New Zealand Trichocolea (Hepaticae), 78
recondensation, in sporogenesis, 145
reduced branches, in Barbula torquata, 483; in Dicranum flagellare, 483; in Tortella calycina, 483
REESE, W. D. (review), H. NOWAK, Revision der Laubmoosgattung Mitthyridium (Mitten) Robinson für Ozeanien (Calymperaceae), 613
REESE, W. D. *See* PURSELL, R. A. & W. D. REESE, 526, 559
REESE, W. D. & I. R. BERMÚDEZ R. Some new records of mosses for Venezuela, 250
regeneration, in Dawsonia leaves, 542+, 543*; in Ditrichum pallidum, 311; in Polytrichum, 309; in Sphagnum, 309
relative cover, bryophytes in sinks, 162+; frequency, bryophytes in sinks, 162+
Relicina, in India, 77, in Sri Lanka, 77; abstrusa, 77+; amphithrix, 77+; relicinula, 78; substrusa, 77+; subconnivens, 78; sublaena, 78; sydneyensis, 78
REYNOLDS, D. Gamete dispersal in Mnium ciliare, 73
Rhabdoweisia crispata, 238
Rhacocarpaceae, 474
Rhacomitrium heterostichum, 254; lanuginosum, 84
Rhacopilum tomentosum, 557
Rhizocarpon badioatrum, 92; chionophilum, 92;

disporum, 92; ferax, 90+; geographicum, 92; lecanorinum, 91
Rhizogoniaceae, 474
Rhizogonium mniodes, 474; spiniforme, 557
rhizoid production, effect of stream flow, 477; effect of temperature, 477
rhizoids, acrocarpous mosses, 481; pleurocarpous mosses, 481
Rhodobryum roseum, 165+, 207
Rhynchostegium serrulatum, 207, 307
Rhytiadelphus squarrosus, 238; triquetrus, 73, 238, sensitivity to trampling, 302
Rhytidium rugosum, 238, 362
Riccardia chamedryfolia, 100; latifrons, 100; multifida, 238, var. multifida, 56, 100, 354; palmata, 100; pinguis, sporogenesis in, 159
Riccia, effect of exogenous auxin and cytokinin, 67; duplex, 67+, 69*; fluitans, 70, carotenoids, 23; rhenana, 67+, 69*
Ricciocarpus natans, 100
ring stage, in telophase I in sporogenesis, 155*, 156
Rinodina papillata, 289+; turfacea, 89+
Robergea pupula, 17
Rocky Mountain National Park, Hepaticae in, 351
ROGERS, A. F. *See* NEUMANN, A. J. & A. F. ROGERS, 369
Ross Lake Natl. Rec. Areas, Hepaticae of, 95
RUDOLPH, E. D., 64
RUNDEL, P. W. Corticolous lichen communities of Nothofagus dombeyi on Volcán Villarica in southern Chile, 82
RUSHING, A. E. (review), A. JAEGER & F. W. SAUERBECK, Genera et species muscorum systematicae disposita seu Adumbratio florae muscorum totius terrarum accedunt Musci cleistocarpi, Enumeratio Fissidentacearum, 612
Russula emetica, 330

Sagedia affinis, 18
saxicolous bryophytes 165; lichens, 329
Scabridens sinensis, 454*
Scapania, 40, 497, western North America, 40; subg. Buchiella, 43; subg. Jensenia, 42; subg. Kaalaasia, 42; subg. Protoscapania, 57; subg. Scapania sect. Aequilobae, 57; subg. Umbrosae, 57; subg. Scapaniella, 43; sect. Apiculatae, 43; sect. Brevicaulae, 45; sect. Curtae, 45; sect. Irriguae, 48; sect. Nemorosae, 55; sect. Scapania subsect. Obscurae, 50, subsect. Scapaniae, 50; americana, 41, 55, 58, 100; apiculata, 41+; bolanderi, 41, 55, 58, 100, 502; carinthiaca, 59; compacta, 58; crassiretis, 41+; curta, 41+, 351+, var. curta, 101, 354; cuspiduligera, 41+, var. cuspiduligera, 354; degenii, 41+, var. dubia, 59; **fulfordiae**, 41, 44*, 58, 354; glaucocephala, 41+; gymnostomophila, 41+; helvetica, 58; hians subsp. salishensis, 59; **hollandiae**, 41+, 53*, 54*; hyperborea, 41+; irrigua, 41+, 351+, var. rufescens, 48, 58; kaurinii, 59; massalongi, 41+; mucronata subsp. mucronata, 41+, subsp. praetervisa, 41+; nemorosa, 59, 307; ornithopodioides, 40+; paludicola, 41+; paludosa, 42+, 101; parvifolia, 41+; saxicola, 43, 59; scandica, 41+; simmonsii, 41+; spitbergensis, 40+; subalpina, 101, 354, 503, var. muddiae, 42+, 49*, 52*, var. subalpina, 42+; uliginosa, 42+, 101; umbrosa, 41+, 102, 238; undulata, 351+, var. oakesii, 51+, 101, 351+, var. undulata, 42+, 101
Scapaniaceae, 497
scariosus-type peristome, 315
Schistostega pennata, 359
Schlotheimia, 405+; tecta, 557; gracillima, 433; pulchella, 420*, 424; torquata, 557
Schoenobryum gardneri, 557
Schofieldia monticola, 101, 226
Schwetschkeopsis denticulata, 168, 207; fabronia, 165
Sciaromnium fryei, 521+; lescurii, 207, 316; tricostatum, 521+, 522*, in Hawaiian Islands, 521, in Oregon, 521, taxonomic affinities, 523
Scorpidium scorpioides, 492
selectivity coefficient, lichens, 35
Selegeria, 559; donniana, 362; pusilla, 167+; tristichoides, 359
SELKIRK, P. M. Effect of an exogenous auxin and cytokinin on Riccia, 67
SELKIRK, P. M. Regeneration from Dawsonia leaves, 542
Sematophyllum adnatum, 207; carolinianum, 306; demissum, 207
SEPPELT, R. D. What is Ditrichum falcifolium? 589
septal formation, in mosses, 159
sexual reproduction, energy investment in, 344; reproductive patterns, in Parmotrema, 344
shade plants, 85
shales, Ordovician calcareous, 338
SHARP, A. J. *See* PETERSON, W. L. & A. J. SHARP, 370
SHARP, A. J. *See* SMITH, D. K. & A. J. SHARP, 401
shoot buds, 310
SHOWERS, D. W. Mielichhofera **tehamensis**, sp. nov., from Northern California, 365
SHUSHAN, S., ABLS representative to AAAS, 129
"shuttle-strategy," 349
SINGH, K. P. A new species of Parmelina from India, 533
sinks, microhabitats in, 165; moisture levels in, 165; temperature in, 165; vapor pressure deficits in, 165

SMIRNOVA, Z. N. (death), 279
SMITH, D. H. Funaria americana Lindb. in North America, 315
SMITH, D. K. & A. J. SHARP (review), H. S. CONARD & P. L. REDFEARN, JR., How to know the mosses and liverworts, 401
SNELGAR, W. P. & T. G. A. GREEN. Carbon dioxide exchange in lichens: low carbon dioxide compensation levels and lack of apparent photorespiratory activity in some lichens, 505
SNIDER, J. A. *See* LODWICK, L. N. & J. A. SNIDER, 214
Solorina crocea, 90
Sørenson similarity index, 284
SPENCE, E. J., 335
Sphaerophorus fragilis, 92; globosus, 92
Sphagnum, 90, 280, in Texas, 247, regeneration in, 309; **andersonianum**, 60⁺, 61*; angustifolium, 62; annulatum, 63; apiculatum, carotenoids, 24⁺; centrale, 241; cuspidatum, 62, 216; cyclophyllum, 216, 248; fallax, 62; fitzgeraldii, 216; fimbriatum, 62; flexuosum, 62; fuscum, 60⁺; heneryense, 239⁺, 240*, distribution of, 241; imbricatum, 216, 240⁺, 248⁺, var. affine, 216; lescurii, 216⁺, 248, chlorophyll a/b ratio, 85; lindbergii, 60; macrophyllum, 216⁺, 248; magellanicum, 62, 217, 309, carotenoids, 24⁺; majus, 62, 238; molle, 62, new to Texas, 248; nemoreum, 62, carotenoids, 27; palustre, 217, 239⁺, 248⁺; papillosum, 240⁺; perichaetiale, 214⁺, 248; potoricense, 216; pulchricoma, 217; pulchrum, 238; pylaesii, 238; squarrosum, 238; recurvum, 217, new to Texas, 248, sensitivity to trampling, 301, var. amblyphyllum, 217; rubellum, 60⁺, sensitivity to trampling, 301; russowii, 62; splendens, 63; strictum, 216, 240; subsecundum, 207, 217, 249, var. rufescens, 217; tenellum, 60⁺; tenuifolium, 60; torreyanum, 216; trinitense, 216
spindle structure, in sporogenesis, 158
Splachnum, 280
splash cups, 74
spore germination, in Jubula, 197⁺; separations, in mosses, 159
spores, achlorophyllous, 554; chlorophyllous, 554; in Leptodictyum riparium, 545⁺
sporocyte, lipid bodies in, 138⁺; oil bodies in, 141
sporogenesis, anaphase I in, 155*; cleavage patterns in, 159; in Ditrichum pallidum, 137⁺, 153⁺; in Hypnum rusciforme, 159; in Riccardia pinguis, 159; meiotic prophase, 137⁺; metaphase I in Ditrichum pallidum, 153⁺, 154*; microtubules in, 153⁺, 154*; mitochondria in, 143, 155; nuclear envelope in, 153; phragmoplast in, 157; spindle structure in, 158; telophase I, 156, ring stage in, 155*, 156; tetrad in Ditrichum pallidum, 153⁺; ultrastructure, 137⁺, 139*, 153⁺, 154*
sporophyte anatomy, in Jubula, 195⁺; development, in Jubula, 195
squamitic acid, 342
Stansted cell disrupter, 342
starch grains, in moss spores, 547
STARK, L. R. Triquetrella in North America, 363
stem formation, in Jubula, 191⁺
Stereocaulon alpinum, 88⁺; arenarium, 90⁺; articum, 90; ramulosum, CO_2 compensation level in, 506; speciosum, 84; testudinea, 92; vesuvianum, 90⁺
Stereodon, 460
Stereophyllum cultelliforme, 557
Sticta filix, CO_2 compensation level in, 506; fuliginosa, 30, 37, 220, CO_2 compensation level in, 506; hypochra, 83; latifrons, CO_2 compensation level in, 506; weigelii, 30⁺
stictic acid, 331⁺
Stokesiella praelonga, 521
stoloniferous branches, as propagules, 483
STONEBURNER, A. *See* WYATT, D., D. M. LANE & A. STONEBURNER, 247
STONEBURNER, A. *See* WYATT, R. & A. STONEBURNER, 512
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature on hepatics and anthocerotes—86, 111; —87, 268; —88, 373; —89, 592
stress adaptation, in bryophytes, 349; in lichens, 349
Strigula, 17; **affinis**, 18; **americana**, 18; complanata, 18; elegans, 18; **phaea**, 18; **viridiseda**, 18
STUDLAR, S. M. Trampling effects on bryophytes: trail surveys and experiments, 301
subbasilaris-type peristome, 323
sulfur-dioxide, bioassay, 281; effects on lichen communities, 281⁺; effects on morphology of lichens, 296⁺
SULLIVANT, W. S., 335
synaptinemal complexes, 142⁺
synchrony, in meiosis, 145
syncytium, 145
Syrrhopodon annotinus, 250; circinatus, 250; ligulatus, 250; parasiticus, 557; prolifera var. circinnatus, 251, var. papillosus, 557, var. prolifer, 251, tenuifolius, 251, 557

TABER, W. A. (review), JAN KOHLMEYER & ERIKA KOHLMEYER, Marine Mycology. The Higher Fungi, 127
Tasmania, Hepaticae in, 220
taxifolius-type peristome, 323
Taxiphyllum, 461⁺; deplanatum, 165⁺, 207; mexicanum, 463; planissimum, 557; scalpellifolium, 557; taxirameum, 165⁺, 207
Tayloria lingulata, 238, 359
telophase I, in sporogenesis, 156
Teloschistaceae, 64
Teloschistes chrysophthalmus var. cinereus, 220; exilis, 220

temperature, in sinks, 165
Tennessee, limestone sinks in, 161+
Tetraphis, 280
Tetraplodon pallidus, 359
Texas, mosses of, 278; Sphagnum in, 214
Thamnobryum alleghaniense, 165+
Thamnolia subuliformis, 88+
Thamnoma constipans, 65
Thelenella amylospora, 7
Thelia asprella, 207; chlorophyll a/b ratio, 85; hirtella, 207; lescurii, 207
Thelopsis inordinata, 18; rubella, 19
Thelotrema hians, 368; **neohians**, 368
Thuidium delicatulum, 165+, 207, chlorophyll a/b ratio, 85, 86, sensitivity to trampling, 301; involvens, 557; minutulum, 207; pygmaeum, 165+, 207; recognitum, 207; urecolatum, 558
thylacoids, in moss spores, 547
Timmia austriaca, 238, subfossils, 487; norvegica var. norvegica, 359
Tomasellia **americana**, 19; **eschweileri**, 19; lactea, 19
Tomenthypnum nitens, 457, 492+
Tortella calycina, reduced branches in, 483; humilis, 165+, 207; inclinata, fossils, 486; richardsii, 558; tortuosa, 207+, 558
Tortula amplexa, 210; bistratosa, 492; cainii, 209+; fragilis, 207; latifolia, 210; mucronifolia, 359, 362; muralis, 207; norvegica, 209+; pagorum, 207; ruralis, 207, 238, var. crinita, 211
trabeculare, 315+
Trachyphyllum, 460
trail survey, sampling methods, 302
trampling, effects in acrocarpous mosses, 311; effects on pleurocarpous mosses, 311; resistant structural features, 311
transect, line-intercept, 162
Trebouxia, 175
Trematodon boasii, 357
Trentepohlia, 6, 170+; aurea, 175
Trichocolea, in New Zealand, 78, in Tasmania, 78; australis, 79; elegans, 79; hatcheri, 79+; julacea, 81; lanata, 81; mollissima, 79+; novaezelandiae, 79; rigida, 79+ tomentella, 238
Trichosteleum popkvilense, 277; sublongisetum, 277
Trichostomeae, European, 132
Trichostomum binnsii, 367; jamaicense, 207; tenuirostris, 165+
Trichothelium epiphyllum, 19; horridulum, 19
Triquetrella, 474; californica, 230, 363; ferruginea, 230, 363
Tritomaria exsecta, 45; exsectiformis, 351+, var. exsectiformis, 101; scitula, 101
Trypethelium, 7; eluteriae, ultrastructure in, 170+, 172*
TUCKER, S. C. New and noteworthy pyrenocarpous lichens from Louisiana and Florida, 1
TUCKER, S. C. *See* LAMBRIGHT, D. D. & S. C. TUCKER, 170
Tuerckheimia linearis, 558

Ulota crispa, 238; drummondii, 238; phyllantha, 238; ulophylla, carotenoids, 24, 25
ultrastructure, chloroplast, 545+, 546*–552*; in Trypethelium eluteriae, 170
Umbilicaria arctica, 90, 92; cylindrica, 92; hyperborea, 92; torrefacta, 92
Usnea, 175; herrei, 330; hirta, 289
usnic acid, 243+, 331, 342

vapor pressure deficits in sinks, 165
VECKER, V. E. Nitrogen fixing lichens of the southern Appalachian mountains of North Carolina, 29
Venturiella sinensis var. angustiannulata, 207
Verrucaria contendens, 12; concatervans, 15; phaea, 18; prospersella, 13; quinqueseptata, 12; stigmatella var. lactea, 19; subcinerea, 12; subprostrans, 4; tamarindi, 4; viridiseda, 18
violaxanthin, 22+
VITT, D. H. The genus Macrocoma. I. Typification of names and taxonomy of the species, 405
VITT, D. H. The genus Macrocoma. II. Geographical variation in the Macrocoma tenue-M. sullivantii species complex, 437
VITT, D. H. & W. J. HOE. Andreaea in Hawaii, 212
vittae, 498

Wabash hydraulic bomb, 343
WEBER, W. A. (review), G. SCHNEIDER, Die Flechtengattung Psora sensu Zahlbruckner, 611
Weissia andrewsii, 207; controversa, 165, 207, 238, 545
Wenatchee Natl. Forest, Hepaticae of, 95
WETMORE, C. M. A new species of Nephroma from North America, 243
Weymouthia mollis, 84
WILL-WOLF, S. Effects of a "clean" coal-fired power plant generating station on four common Wisconsin lichen species, 296
WILL-WOLF, S. Structure of corticolous lichen communities before and after exposure to emissions from a "clean" coal-fired generating station, 281
WYATT, R., D. M. LANE & A. STONEBURNER. Additions to the Sphagnum flora of Texas, 247
WYATT, R. & A. STONEBURNER. Distribution and phenetic affinities of Donrichardsia, an endemic moss from the Edwards Plateau of Texas, 512

Xanthoparmelia cumberlandia, 330+
Xanthoria candelaria, 90+; elegans, 90+; fallax, 220, 289; parietina, 220, 505
Xylographa micrographa, 71

ZANDER, R. H. Acid-base color reactions: The status of Triquetrella ferruginea, Barbula inaequalifolia, and B. calcarea, 228
ZANDER, R. H., Editorial, 617
ZANDER, R. H. *See* BUCK, W. R. & R. H. ZANDER, 254
ZANDER, R. H. *See* JANSSENS, J. A. & R. H. ZANDER, 486
ZANDER, R. H., C. DELGADILLO M. & P. M. ECKEL. Morinia saitoana, a new species of Pottiaceae (Bryopsida) from Mexico, 508
ZANDER, R. H. & P. M. ECKEL. Tortula cainii: additional Ontario records and behavior in a common garden, 209
α-zeacarotene, 26; β-, 26
zeaxanthin, 22+
zeorine, 243
Zygodon pentastichus, 84

THE BRYOLOGIST

Quarterly Journal of the American Bryological and Lichenological Society

Editor
RICHARD H. ZANDER

Volume 83
1980

Table of Contents

ALLEN, B. H. Mosses new or noteworthy for Colorado 257
ALLEN, B. H. Peristome variations in the Genus *Fissidens*: an SEM study 314
ANDERSON, L. E. Review 394
ANDRUS, R. E. *Sphagnum andersonianum*, a new species endemic to northeastern North America 60
ANDRUS, R. E. A large range extension for *Sphagnum henryense* 239
BECKER, V. E. Nitrogen fixing lichens in forests of the Southern Appalachian Mountains of North Carolina 29
BELLAND, R. J. & G. R. BRASSARD. New or additional moss records from Newfoundland VI 360
BELLAND, R. J. & G. R. BRASSARD. A major range extension for *Encalypta longicolla* 563
BRAGG, T. B. Review 401
BRASSARD, G. R. Bryophyta Exsiccata Terrae-Novae et Labradoricae: a new exsiccata from Newfoundland and Labrador, Canada 237
BRODO, I. M. Review 614
BROWN, R. C. & B. E. LEMMON. Ultrastructure of sporogenesis in a moss, *Ditrichum pallidum* I. Meiotic propase 137
BROWN, R. C. & B. E. LEMMON. Ultrastructure of sporogenesis in a moss, *Ditrichum pallidum* II. Metaphase I through the tetrad 153
BROWN, R. C. & B. E. LEMMON. Ultrastructural aspects of chloroplast development in spores of the moss *Leptodictyum riparium* (Hedw.) Warnst. 545
BRYSON, C. T. *Diphyscium foliosum* new to Mississippi 372
BUCK, W. R. Review 278
BUCK, W. R. Animadversions on *Pterigynandrum* with special commentary on *Forsstroemia* and *Leptopterigynandrum* 451
BUCK, W. R. & R. H. ZANDER. On *Astomiopsis*, *Bryomanginia* and *Melophyllum* (Ditrichaceae) 254
CHRISTY, J. A. Additions to the moss flora of Oregon 355
CHRISTY, J. A. Rediscovery of *Sciaromium tricostatum* (Sull.) Mitt. (= *Limbella tricostata* (Sull.) Bartr.) in Oregon 521
CRANDALL-STOTLER, B. & W. R. GUERKE. Developmental anatomy of *Jubula* Dum. (Hepaticae) 179
CROSBY, M. R. Recent literature on mosses—114–116 ..... 118, 270, 388
CROSBY, M. R. & C. R. BAUER. Recent literature on mosses—117 603
CULBERSON, W. L. Review 403
CZECZUGA, B. Investigations on carotenoids in Embryophyta. I. Bryophyta 21
DURY, M. N. Maurice Louis Jean Bizot (1905–1979) 564
EGAN, R. S. Recent literature on lichens—104–107 ..... 103, 263, 380, 596
EGAN, R. S. Reviews 613, 614
ENGEL, J. J. Austral hepaticae XII. A new species of *Clasmatocolea* (Hepaticae) from Tasmania 220
ESSLINGER, T. L. Typification of *Oropogon loxensis* and description of two related species 529
FAHSELT, D. Alternative method for analyzing protein characters in lichens 340
FIFE, A. J. The affinities of *Costesia* and *Neosharpiella* and notes on the Gigaspermaceae (Musci) 466

FIFE, A. J. & G. R. BRASSARD. New or additional moss records from Newfoundland V 358
FRAHM, J.-P. Review 277
FRAHM, J.-P. Synopsis of the genus *Campylopus* in North America north of Mexico 570
GLIME, J. M. Effects of temperature and flow on rhizoid production in *Fontinalis* 477
GODFREY, J. D. & G. A. GODFREY. Notes on hepatics from the Pacific Northwest 224
HALE, M. E., JR. The lichen genus *Relicina* (Parmeliaceae) in India and Sri Lanka 77
HANSEN, E. S. Lichens from northwestern Greenland collected on botanical expeditions in 1975 and 1977 87
HARPEL, J. S. *Orthotrichum pumilum* new to California 201
HATTAWAY, R. A. The calciphilous bryophytes of three limestone sinks in eastern Tennessee 161
HERMANN, F. J. Distinctions between *Anacolia menziesii* (new to Colorado) and *Bartramia stricta* 253
HERRNSTADT, I., C. C. HEYN & M. R. CROSBY. New data on the moss genus *Gigaspermum* 536
HONG, W. S. The genus *Scapania* in western North America. II. Taxonomic treatment 40
HONG, W. S. Hepaticae of the North Cascades Range, Washington 94
HONG, W. S. Hepaticae of Rocky Mountain National Park, Colorado 351
HONG, W. S. A study of the distribution of *Diplophyllum* in western North America 497
IRELAND, R. R. The taxonomic status of *Ceratodon heterophyllus* 234
JACKSON, D. W. & A. W. HOPKINS. New additions to the lichen flora of Texas and the United States 532
JANSSENS, J. A. & R. H. ZANDER. *Leptodontium flexifolium* and *Pseudocrossidium revolutum* as 60,000-year-old subfossils from the Yukon Territory, Canada 486
JORDAN, W. P. *Edrudia*, a new genus from California (Lichens: Teloschistaceae) .. 64
JUDD, W. S. Notes on the moss flora of Hispaniola 555
KEENER, C. S. Review 615
LAMBRIGHT, D. D. & S. C. TUCKER. Observations on the ultrastructure of *Trypethelium eluteriae* Spreng. 170
LAWREY, J. D. Correlations between lichen secondary chemistry and grazing activity by *Pallifera varia* 328
LAWREY, J. D. Sexual and asexual reproductive patterns in *Parmotrema* (Parmeliaceae) that correlate with latitude 344
LESQUEREUX, L. Letter 128
LODWICK, L. N. & J. A. SNIDER. The distribution of *Sphagnum* taxa in Texas 214
LORIA, M. & I. HERRNSTADT. Moss capsules as food of the harvester ant, *Messor* .. 524
MAHLER, B. D. & W. F. MAHLER. Checklist of mosses of Oklahoma 202
MARTIN, C. E. Chlorophyll a/b ratios of eleven North Carolina mosses 84
MOBERG, R. *Anaptychia ulotrichoides* new to North America 251
NEUMANN, A. J. & A. F. ROGERS. Contribution to the bryoflora of Texas 369
NOBLE, W. J. *Catillaria columbiana* comb. nov. from western North America ... 71
OSORIO, H. S. Contribution to the lichen flora of Argentina XII. New or additional records from Buenos Aires Province 219

PATWARDHAN, P. G. & U. MAKHIJA. Nomenclatural note on three species of *Anthracothecium* 368
PECK, J. H. *Bryoxiphium norvegicum* in Iowa 535
PETERSON, W. & A. J. SHARP. *Leucobryum incurvifolium* in the United States 370
PURSELL, R. A. *Fissidens diplodus* Mitt. var. *richardsii* (R. S. Williams) Pursell, comb. nov. 256
PURSELL, R. A. & W. D. REESE. The rediscovery of *Fissidens subulatus* Mitt. in Brazil 526
PURSELL, R. A. & W. D. REESE. The taxonomic status of the Nanobryaceae (Bryopsida) 559
RATKOWSKY, D. A. Tasmanian and New Zealand *Trichocolea* (Hepaticae) 78
RATKOWSKY, D. A. Some further synonyms of *Bryoerythrophyllum jamesonii* (Tayl.) Crum 367
REESE, W. D. Review 613
REESE, W. D. & I. R. BERMÚDEZ. Some new records of mosses for Venezuela 250
Reynolds, D. N. Gamete dispersal in *Mnium ciliare* 73
RUNDEL, P. W. Corticolous lichen communities of *Nothofagus dombeyi* on Volcán Villarica in southern Chile 82
RUSHING, A. E. Review 612
SMITH, D. K. & A. J. SHARP. Review 401
SELKIRK, P. M. Effect of an exogenous auxin and cytokinin on *Riccia* 67
SELKIRK, P. M. Regeneration from *Dawsonia* leaves 542
SEPPELT, R. D. What is *Ditrichum falcifolium*? 589
SHOWERS, D. W. *Mielichhoferia tehamensis*, sp. nov., from northern California . 365
SINGH, PAL, KRISHNA. A new species of *Parmelina* from India 533
SMITH, D. K. *Funaria americana* Lindb. in North America 335
SNELGAR, W. P. & T. G. A. GREEN. Carbon dioxide exchange in lichens: low carbon dioxide compensation levels and lack of apparent photorespiratory activity in some lichens 505
STARK, L. R. *Triquetrella* in North America 363
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature on hepatics and anthocerotes—86–89 ..... 111, 258, 373, 592
STUDLAR, S. M. Trampling effects on bryophytes: trail surveys and experiments .. 301
TABER, W. A. Review 128
TUCKER, S. C. & R. C. HARRIS. New and noteworthy pyrenocarpous lichens from Louisiana and Florida 1
VITT, D. H. The genus *Macrocoma*. I. Typification of names and taxonomy of the species 405
VITT, D. H. The genus *Macrocoma*. II. Geographical variation in the *Macrocoma tenue-M. sullivantii* species complex 437
VITT, D. H. & W. J. HOE. *Andreaea* in Hawaii 212
WEBER, W. A. Review 611
WETMORE, C. M. A new species of *Nephroma* from North America 243
WILL-WOLF, S. Structure of corticolous lichen communities before and after exposure to emissions from a "clean" coal-fired generating station 281
WILL-WOLF, S. Effects of a "clean" coal-fired power generating station on four common Wisconsin lichen species 296
WYATT, R. & A. STONEBURNER. Distribution and phenetic affinities of *Donrichardsia*, an endemic moss from the Edwards Plateau of Texas 512

WYATT, R., D. M. LANE & A. STONEBURNER. Additions to the *Sphagnum* flora of Texas 247
ZANDER, R. H. Editorials 133, 617
ZANDER, R. H. Acid-base color reactions: the status of *Triquetrella ferruginea*, *Barbula inaequalifolia* and *B. calcarea* 228
ZANDER, R. H. & P. M. ECKEL. *Tortula cainii*: additional Ontario records and behavior in a common garden 209
ZANDER, R. H., C. DELGADILLO M. & P. M. ECKEL. *Morinia saitoana*, a new species of Pottiaceae (Bryopsida) from Mexico 508

Actual dates of issue of Volume 83:
Number 1: 17 March 1980
Number 2: 13 August 1980
Number 3: 21 November 1980
Number 4: 9 June 1981